

Dirac oscillator in an external magnetic field

Bhabani Prasad Mandal* and Shweta Verma†

Department of Physics, Banaras Hindu University, Varanasi-221005, India

We show that 2+1 dimensional Dirac oscillators in an external magnetic field is mapped onto the same with reduced angular frequency in absence of magnetic field. This can be used to study the atomic transitions in a radiation field. Relativistic Landau levels are constructed explicitly. Several interesting features of this system are discussed.

PACS numbers: 31.30.J-; 03.65.Pm

Keywords: Dirac Oscillator, Jaynes-Cumming Model, Relativistic Landau Levels, Graphene

* Email: bhabani.mandal@gmail.com; Phone No: 91-542-6701545

† Present Address: Raja Ramanna Centre for Advanced Technology, Indore-452013

Dirac oscillator [1] was introduced for the first time by Moshinsky and Szczepaniak by adding the linear term $-imc\omega\beta\boldsymbol{\alpha} \cdot \mathbf{r}$ to the usual Dirac Hamiltonian for a free particle and described by the Hamiltonian,

$$H = c\boldsymbol{\alpha} \cdot (\mathbf{p} - im\omega\beta\mathbf{r}) + \beta mc^2, \quad (1)$$

where $\boldsymbol{\alpha}$, β are usual Dirac matrices, m is the rest mass of the particle and c is the speed of light. In the nonrelativistic limit, it reduces to a quantum simple harmonic oscillator (with mass m , angular frequency, ω) with a strong spin-orbit coupling. The Dirac oscillator has attracted a lot of attention and has found many physical applications in various branches of physics [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18]. Initially it was introduced in the context of many body theory, mainly in connection with quark confinement models in quantum chromodynamics (QCD) and in the context of relativistic quantum mechanics [3]. Later the subsequent studies have revealed several exciting properties connected to the symmetries of the theory [11, 12]. Most exciting properties of Dirac oscillator is its connection to quantum optics [10, 11]. In 2+1 dimensions it maps into Anti-Jaynes-Cummings (AJC) model [19, 20, 21] which describes the atomic transitions in a two level system.

In this present article we have studied Dirac oscillator in presence of external magnetic field. The dynamics of the Dirac oscillator in the external uniform magnetic field, $\mathbf{B}$ is governed by the Hamiltonian,

$$H = c\boldsymbol{\alpha} \cdot (\mathbf{p} - \frac{e\mathbf{A}}{c} - im\omega\beta\mathbf{r}) + \beta mc^2. \quad (2)$$

$\mathbf{A}$ is the vector potential which produces the external magnetic field and e is the charge of the Dirac oscillator which is considered to be an electron in this work. This theory of 2+1 dimensional Dirac oscillators in presence of external transverse magnetic field is mapped onto the theory of Dirac oscillators without magnetic field in the same space-time dimension and described by the Hamiltonian,

$$H' = c\boldsymbol{\alpha} \cdot (\mathbf{p} - im\tilde{\omega}\beta\mathbf{r}) + \beta mc^2. \quad (3)$$

The external magnetic field causes the angular frequency of the Dirac oscillator to change by half of the cyclotron frequency. We have solved 2+1 dimensional Dirac equation corresponding to Hamiltonian in Eq. (2) for Dirac oscillator in the presence of external, constant magnetic field exactly and have obtained energy eigenvalues and eigen functions analytically.

Relativistic Landau levels [22] are constructed explicitly. Infinitely degenerate lowest Landau level are constructed in the coordinate basis in a very simple manner. Further, we derive exact mapping between this relativistic model and AJC model so widely used in quantum optics. This relativistic system has many peculiar properties some of which are similar to that of graphene [23, 24, 25, 26, 27], monolayer of carbon atoms.

For the sake of simplicity, we consider the magnetic field, $\mathbf{B}$ along the z -direction. The vector potential for this particular magnetic field can be chosen in the symmetric gauge as $\mathbf{A} = (-\frac{B}{2}y, \frac{B}{2}x, 0)$. Further we are interested to study the system in two space and one time dimension where Dirac matrices can be chosen in terms of 2×2 Pauli spin matrices. We consider one of the widely used choices of Dirac matrices as $\alpha_x = \sigma_x, \alpha_y = \sigma_y$ and $\beta = \sigma_z$. The Hamiltonian in Eq.(2) can be written nicely in terms of complex coordinate $z = x + iy$ and conjugate momentum, p_z , as

$$H = \begin{pmatrix} mc^2 & 2cp_z + im\tilde{\omega}c\bar{z} \\ 2cp_{\bar{z}} - im\tilde{\omega}cz & -mc^2 \end{pmatrix}, \quad (4)$$

with

$$p_z = -i\hbar \frac{d}{dz} = \frac{1}{2}(p_x - ip_y),$$

$$p_{\bar{z}} = -i\hbar \frac{d}{d\bar{z}} = \frac{1}{2}(p_x + ip_y), \quad \text{and} \quad (5)$$

$$\tilde{\omega} = \omega - \frac{\omega_c}{2} \quad \text{where } \omega_c = \frac{|e|B}{mc}, \quad (6)$$

ω_c is the cyclotron frequency. These complex coordinates and their canonical conjugate momenta satisfy the following commutation relations, i.e.

$$[z, p_z] = i\hbar = [\bar{z}, p_{\bar{z}}], \quad ; \quad [\bar{z}, p_z] = 0 = [z, p_{\bar{z}}]. \quad (7)$$

We define the following creation and annihilation operators as

$$a = \frac{1}{\sqrt{m\tilde{\omega}\hbar}}p_{\bar{z}} - \frac{i}{2}\sqrt{\frac{m\tilde{\omega}}{\hbar}}z,$$

$$a^\dagger = \frac{1}{\sqrt{m\tilde{\omega}\hbar}}p_z + \frac{i}{2}\sqrt{\frac{m\tilde{\omega}}{\hbar}}\bar{z}. \quad (8)$$

It can be verified using Eqs. (7) and (8) that $[a, a^\dagger] = 1$; $[a, a] = 0 = [a^\dagger, a^\dagger]$.

The Hamiltonian in Eq.(4) further is expressed in terms of creation and annihilation operators as

$$H = \begin{pmatrix} mc^2 & 2c\sqrt{m\tilde{\omega}\hbar} a^\dagger \\ 2c\sqrt{m\tilde{\omega}\hbar} a & -mc^2 \end{pmatrix}. \quad (9)$$

The complete solution of the Dirac oscillator in the transverse magnetic field is obtained by solving the time independent Dirac equation, $H\Psi = E\Psi$ corresponding to the above Hamiltonian. The relativistic Landau levels are given as,

$$E_n^\pm = \pm mc^2 \sqrt{1 + \frac{4\hbar\tilde{\omega}}{mc^2}n}; \quad n = 0, 1, 2 \dots \quad (10)$$

with normalized negative and positive energy states are given by

$$|\Psi_n^\pm\rangle = c_n^\pm |n; \frac{1}{2}\rangle + d_n^\pm |n-1; -\frac{1}{2}\rangle, \quad (11)$$

with $c_n^\pm = \pm \sqrt{\frac{E_n^\pm \pm mc^2}{2E_n^\pm}}$ and $d_n^\pm = \mp \sqrt{\frac{E_n^\pm \mp mc^2}{2E_n^\pm}}$. We adopt the notation for the state as $|n, \frac{1}{2}m_s\rangle \equiv \psi_n(z, \bar{z})\chi_{m_s}$ where n is the eigenvalue for the number operator, $a^\dagger a$ and $m_s = \pm 1$ are the eigenvalues of the operator σ_z i.e. $\sigma_z|\frac{1}{2}m_s\rangle = m_s|\frac{1}{2}m_s\rangle$, and $\psi_n(z, \bar{z})$ is the space part of the wave function in the coordinate representation whereas χ_{m_s} is the spin part of the wave function. These states indicate the entanglement between orbital and spin degrees of freedom in the Dirac oscillator problem.

The lowest Landau levels [LLL] are obtained by using the condition,

$$a\psi_0(z, \bar{z}) = 0. \quad (12)$$

Equivalently in coordinate representation, the space part wave function ($\psi_0(z, \bar{z})$) of the LLL satisfies

$$\left(\frac{\partial}{\partial \bar{z}} + \frac{m\tilde{\omega}}{2\hbar}z\right)\psi_0(z, \bar{z}) = 0. \quad (13)$$

By substituting

$$\psi(z, \bar{z}) = e^{-\frac{m\tilde{\omega}}{2\hbar}z\bar{z}}u_0(z, \bar{z}), \quad (14)$$

we further obtain

$$\frac{\partial}{\partial \bar{z}}u_0(z, \bar{z}) = 0, \quad (15)$$

as the defining rule for the LLL. We obtain the LLL in the coordinate representation, which is infinitely degenerate, as

$$\psi_0(z, \bar{z}) = z^l e^{-\frac{m\tilde{\omega}}{2\hbar}z\bar{z}}, \quad l = 0, 1, 2 \dots, \quad (16)$$

as $u_0(z, \bar{z})$ is analytic function and monomials z^l with $l = 0, 1, 2 \dots$ can serve as a linearly independent basis. The first excited state and other higher states in the coordinate space can be obtained by applying $a^\dagger$ given in Eq. (8) repeatedly.

We observe several interesting features of this system and study their implications and consequences. Firstly we observe that the 2+1 dimensional Dirac oscillator having angular frequency, ω in the transverse magnetic field maps into a 2+1 dimensional Dirac oscillator in absence of magnetic field but having a different angular frequency, $\tilde{\omega} = \omega - \frac{1}{2}\omega_c$. The effect of transverse magnetic field in the system of 2+1 dimensional Dirac oscillator is to decrease the angular frequency by half of the cyclotron frequency only and the entire dynamics remains unchanged except a special choice on the strength of the magnetic field, $B = \frac{2mc\omega}{|e|}$ for which $\tilde{\omega} = 0$ and Dirac oscillator stops oscillating. This is highly non-trivial situation and the dynamics is completely different from the dynamics of the system without magnetic field.

To show this result we consider the Hamiltonian for Dirac oscillator without magnetic field given in Eq. (1) in 2+1 dimension and express it using complex coordinates and their canonically conjugate momenta listed in Eq. (5) as

$$H = \begin{pmatrix} mc^2 & 2cp_z + im\omega c\bar{z} \\ 2cp_{\bar{z}} - im\omega cz & -mc^2 \end{pmatrix} \quad (17)$$

This Hamiltonian is exactly same (except ω is replaced by $\tilde{\omega}$) as the Hamiltonian given in Eq. (4) which describes the Dirac oscillator in the transverse magnetic field in 2+1 dimensions. Dirac oscillator in presence of external fields is also considered by others [4, 5, 6, 8] to discuss different issues, in particular, to study Aharonov-Bohm effect [6], effect of electric field [8]. 2+1 dimensional Dirac oscillator in the transverse magnetic field is also considered in Ref. [4] to discuss different aspect of chiral quantum phase transition in such a system.

Interestingly we note the spectrum of Landau levels in Eq. (10) are non-equidistant in energy and proportional to the $\sqrt{nB}$ for sufficiently large transverse magnetic field. This feature is quite different from the conventional two dimensional electronic system. However this feature is very similar to the surprising material graphene, the sheet of carbon atoms [23, 24, 25, 26, 27]. The peculiar properties of graphene mainly stem from its linear dispersion near Fermi surface, and the chiral nature of electronic states entangling the momentum and pseudo-spin of the quasi-particle arising due to sublattice structure of carbon atoms in graphene. This entanglement of states in graphene is exactly similar to the entanglement of orbital and spin states in Dirac oscillator given in Eq. (11). Relativistic Landau Levels are experimentally realized [29].

Now we would like to show the connection of this model to the AJC model so widely used in the study of quantum optics. The Hamiltonian corresponding to the simple version

of AJC model is given as,

$$H_{AJC} = g(\sigma^- a + \sigma^+ a^\dagger) + \sigma_z mc^2, \quad (18)$$

On the other hand the JC model is described by the Hamiltonian,

$$H_{JC} = g(\sigma^+ a + \sigma^- a^\dagger) + \sigma_z mc^2, \quad (19)$$

which is quite different from Hamiltonian of AJC model as the positions of creation and annihilation operators are interchanged. The Hamiltonian in the Eq. (9) can be expressed exactly in the form of H_{AJC} as given in Eq. (18) using Pauli spin matrices where $\sigma^\pm = \frac{1}{2}(\sigma_x \pm i\sigma_y)$ are usual spin raising and lowering operators and $g = 2c\sqrt{m\tilde{\omega}\hbar} = 2c\sqrt{m\hbar(\omega - \frac{|e|B}{2mc})}$, coupling between spin and orbital degrees of freedom which depends on the strength of the magnetic field. It is very interesting to note how two completely different theories are related. This connection is established for arbitrary strength of the magnetic field hence differs with some earlier work [4]. This provides an alternative approach to study atomic transitions in two level system using relativistic quantum mechanical model, in particular using the theory of Dirac oscillators in the transverse magnetic fields. The connection between relativistic quantum mechanics and quantum optics using trapped ions is realized experimentally in Ref. [30]

We further observe that the zitterbewegung frequency for the 2+1 dimensional Dirac oscillator in the external magnetic field depends on the strength of the magnetic field. To show this we start with some initial pure state at $t = 0$,

$$|n, \frac{1}{2} > (0) = \frac{E_n^+}{mc^2} [d_n^+ |\Psi_n^- > - d_n^- |\Psi_n^+ >]. \quad (20)$$

This equation is obtained from the Eq.(11) by eliminating the state $|n-1; -\frac{1}{2} >$. The Eq. (20) shows that the starting initial state is a superposition of both the positive and negative energy solutions. The time evolution of this state can be written as

$$|n, \frac{1}{2} > (t) = \frac{E_n^+}{mc^2} [d_n^+ |\Psi_n^- > e^{i\omega_n t} - d_n^- |\Psi_n^+ > e^{-i\omega_n t}], \quad (21)$$

where ω_n is the frequency of oscillation between positive and negative energy solutions and given as

$$\omega_n = \frac{E_n^+}{\hbar} = \frac{mc^2}{\hbar} \sqrt{1 + \frac{4\hbar\tilde{\omega}}{mc^2} n} \text{ with } \tilde{\omega} = \omega - \frac{|e|B}{2mc} \quad (22)$$

Substituting Ψ^+ and Ψ^- and the constants d_n^+ and d_n^- in Eq. (21) we obtain,

$$\begin{aligned}
 |n, \frac{1}{2} \rangle (t) = & [\cos(\omega_n t) - \frac{imc^2}{E_n^+} \sin(\omega_n t)] |n, \frac{1}{2} \rangle (0) \\
 & + i \sqrt{\frac{E_n^{+2} - m^2 c^4}{m^2 c^4}} \sin(\omega_n t) |n-1, -\frac{1}{2} \rangle (0)
 \end{aligned}
 \tag{23}$$

This equation shows the oscillatory behavior between the states $|n, \frac{1}{2} \rangle$ and $|n-1, -\frac{1}{2} \rangle$ which is exactly similar to atomic Rabi oscillations [20, 21, 28] occurring in the JC/AJC models. Rabi frequency is given by ω_n given in Eq. (22).

We would like mention that in the limit magnetic field, $B \rightarrow 0$ all these results are consistent with results of Dirac oscillator (with out magnetic field) in Ref. [10]

In conclusion, we have studied 2+1 dimensional Dirac oscillator in the presence of transverse external magnetic field by defining suitable creation and annihilation operators in terms of properly chosen canonical pairs of coordinates and momenta. We have shown that the Dirac oscillator in magnetic field is mapped onto Dirac oscillator without magnetic field but with different angular frequency. The frequency changes by half the cyclotron frequency. The relativistic Landau levels and corresponding eigenstates have been calculated explicitly by solving corresponding Dirac equation. Lowest Landau levels have been constructed exactly in the coordinate space representations which shows the infinite degeneracy of such state in a simple manner. We have observed several interesting features of such system. Some of the features are quite similar to those shown by graphene. In particular, the orbital and spin degrees of freedom are entangled to each other, the Landau levels are non-equidistant in energy and the level separations are proportional to $\sqrt{B}$ for large magnetic field. The connection between AJC model in quantum optics with Dirac oscillator in magnetic field is established in this work in a simple manner without taking any limit on the strength of the magnetic field. This will allow to study atomic transitions in quantum optics using the relativistic quantum mechanical models in presence of magnetic field. The Zitterbewegung frequency has been calculated for the Dirac oscillator in magnetic field and has shown to vary as $\sqrt{nB}$ for sufficiently large magnetic field. This oscillation between the negative and positive energy states is similar to the Rabi oscillation in the two level systems. It will be interesting to study graphene in Dirac oscillator potential in presence of transverse magnetic

field.

-
- [1] M. Moshinsky & A Szczepaniak ; J. Phys. A22, (1989) L817-L819. Similar system was studied much earlier by Ito et.al [3] and Cook[31]. However the term *Dirac Oscillator* was introduced first time by Moshinsky and Szczepaniak [1].
 - [2] R.Lisboa, M. Malheiro, A.S de Castro, P Albreto, M Foilhais, Phys. Rev C 69 (2004) 024319 (1-19).
 - [3] D Ito, K Mori & E Carrieri, Nuovo Cimento 51 A (1967), 1119-1121.
 - [4] A Bermudez, M A Martin Delgado & A Luis, Phys. Rev. A 77 (2008) 063815 (1-13).
 - [5] E. Sadurni, J. Mauricio Torres, & T. H. Seligman, arXiv: 0902.1476.
 - [6] N. Ferkous & A. Bounames, Phys. Lett A 325, 21 (2004).
 - [7] Ahmed Jellal, Abdulaziz D. Alhaidari, Hocine Bahlouli, arXiv:0904.0587.
 - [8] D.A. Kulikov, & R. S.Tutik, physics/0608277, Proceedings of International Conference on Mathematical Methods in Electromagnetic Theory (MMET*06), Kharkiv, 26-28 June (2006) 554-556.
 - [9] Victor M Villalba Phys Rev. A 49,(1994) 586-587.
 - [10] A Bermudez, M. A.Martin-Delgado and E.Solano, Phys. Rev. A 76, (2007) 041801(R) [1-4].
 - [11] J. Benítez, R. P. Martinez y Romero, H. N. Núñez-Yépez and A. L. Salas-Brito, Phys. Rev. Lett. 64,(1990) 1643-1645; Erratum, Phys. Rev. Lett. 65, (1990) 2085-2085.
 - [12] M. Moreno and A. Zentella, J. Phys. A: Math. Gen 22, (1989) L821-L825 .
 - [13] V M Villalba & A Rincon Maggiolo, Eur. Phys. J. B22 (2001), 31-35.
 - [14] C. Quesne & V.M. Tkachuk, J. Phys A38, (2005) 1747-1765 ; M. Mohadesi & B. Mirza,Commun. Theor. Phys. 42, (2004) 664-668 ; A. D. Alhaidari, Found. Phys. Lett 18,(2005) 651-664.
 - [15] Y. Brihaye, & A. Nininahazwe, Mod. Phys. Lett. A20 (2005) 1875-1885.
 - [16] R. de Lima Rodrigues, Phys. Lett. A372 (2008) 2587-2591.
 - [17] R. P. Martinez-y-Romero, H. N. Núñez-Yépez, A. L. Salas-Brito, Eur.J.Phys 16 (1995) 135-141.
 - [18] P. Rozmej & R. Arvieu, J. Phys. A32 (1999), 5367-5382.
 - [19] E T Jaynes& F W Cummings, Proc. IEEE, 51, (1963) 89-109.

- [20] Jonas Larso, Phys. Scr. 76, (2007) 146-160.
- [21] L Allen and J H Eberly, *Optical Resonance and Two Level Atoms* Dover Publications, Mineola, New York, (1987).
- [22] I. I. Rabi, Zeit. f. Physik 49, (1928) 507-511; M. H. Johnson, B. A. Lippmann, Phys. Rev. 76, (1949) 828-832; M. H. Johnson, B. A. Lippmann, Phys. Rev. 77, (1950) 702-705; A. Bermudez, M. A. Martin-Delgado, and E. Solano, Phys. Rev. Lett. 99, (2007) 123602-123605.
- [23] A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov and A. K. Geim, Rev. Mod. Phys. 81 (2009) 109-163.
- [24] M.I. Katsnelson, K.S. Novoselov & A.K. Geim, Nature Phys. 2 (2006) 620-625.
- [25] K. Shizuya, Phys. Rev. B, 77 (2008) 075419 (1-6).
- [26] J. Schliemann, Phys. Rev. B, 78 (2008), 195426 (1-5) .
- [27] J. Schliemann, New J. Phys. 10 (2008), 043024 (1-9).
- [28] I I Rabi, Phys. Rev. 51 (1937) 652-654.
- [29] G. Li, E. Y. Andrei, Nat. Phys. 3, (2007), 623-627.
- [30] R Gerritsma, G. Kirchmair, F. Zahringer, E. Solano, R. Blatt, and C. F. Roos, arXiv: 0909.0674 (2009).
- [31] P A. Cook, Lettere al Nuovo Cimento, 1 (1971), 419-421.